

Patient name: _____

Physician name: _____

Date: _____

Bath Ankylosing Spondylitis Functional Index (BASFI)

It should take you no more than a minute or two to complete this questionnaire. Your responses will help your doctor assess how active your ankylosing spondylitis (AS) is, and will let him or her track changes in your condition over time.

Please mark an X in the box that corresponds to how easy or difficult it was to perform the activity described (your answers should apply to the past week only). Your doctor will add up the scores.

An aid is a piece of equipment which helps you to perform an action or movement.

1. Putting on your socks or tights without help or aids (e.g., sock aid).

0 1 2 3 4 5 6 7 8 9 10

Easy Impossible

SCORE

2. Bending forward from the waist to pick up a pen from the floor without an aid.

0 1 2 3 4 5 6 7 8 9 10

Easy Impossible

3. Reaching up to a high shelf without help or aids (e.g., helping hand).

0 1 2 3 4 5 6 7 8 9 10

Easy Impossible

4. Getting up out of an armless dining room chair without using your hands or any other help.

0 1 2 3 4 5 6 7 8 9 10

Easy Impossible

5. Getting up off the floor without help from lying on your back.

0 1 2 3 4 5 6 7 8 9 10

Easy Impossible

6. Standing unsupported for 10 minutes without discomfort.

0 1 2 3 4 5 6 7 8 9 10

Easy Impossible

7. Climbing 12-15 steps without using a handrail or walking aid, one foot on each step.

0 1 2 3 4 5 6 7 8 9 10

Easy Impossible

8. Looking over your shoulder without turning your body.

0 1 2 3 4 5 6 7 8 9 10

Easy Impossible

9. Doing physically demanding activities (e.g., physiotherapy exercises, gardening or sports).

0 1 2 3 4 5 6 7 8 9 10

Easy Impossible

10. Doing a full day's activities, whether it be at home or at work.

0 1 2 3 4 5 6 7 8 9 10

Easy Impossible

Last BASFI score

Total Q1 to Q10

Absolute change

Total Q1 to Q10 ÷ 10 = BASFI score

% change

- Adapted from Calin A et al.¹

HUMIRA. Power to Fight AS

15 years of clinical experience worldwide combined across all indications²

>1.6
million
patient-years
of exposure²

>670,000
patients
currently treated
worldwide²

71
clinical trials
with over
23,000
patients²

9
years
in Canadian
practice³

7
indications³

Available
in 89
countries²

HUMIRA[®] | 15 of clinical
adalimumab | years experience
worldwide

Indicated in the inflammatory conditions:³

RA

JIA

PsA

AS

Adult
CD

Ped CD

Ps

HUMIRA is indicated for:

- Reducing the signs and symptoms, inducing major clinical response and clinical remission, inhibiting the progression of structural damage and improving physical function in adult patients with moderately to severely active rheumatoid arthritis (RA). Can be used alone or in combination with methotrexate (MTX) or other disease-modifying antirheumatic drugs (DMARDs).

When used as first-line treatment in recently diagnosed patients who have not been previously treated with MTX, HUMIRA should be given in combination with MTX. Can be given as monotherapy in case of intolerance to MTX or when treatment with MTX is contraindicated.

- In combination with MTX, reducing the signs and symptoms of moderately to severely active polyarticular juvenile idiopathic arthritis (JIA) in patients 4 to 17 years of age who have had an inadequate response to one or more DMARDs. Can be used as monotherapy in case of intolerance to MTX or when continued treatment with MTX is not appropriate. HUMIRA has not been studied in children aged less than 4 years.
- Reducing the signs and symptoms in patients with active ankylosing spondylitis (AS) who have had an inadequate response to conventional therapy.
- Reducing the signs and symptoms of active arthritis and inhibiting the progression of structural damage and improving the physical function in adult psoriatic arthritis (PsA) patients. Can be used in combination with MTX in patients who do not respond adequately to MTX alone.
- Reducing the signs and symptoms and inducing and maintaining clinical remission in adult patients with moderately to severely active Crohn's disease (CD) who have had an inadequate response to conventional therapy, including corticosteroids and/or immunosuppressants. HUMIRA is indicated for reducing the signs and symptoms and inducing clinical remission in these patients if they have also lost response to or are intolerant to infliximab.
- Reducing the signs and symptoms and inducing and maintaining clinical remission in pediatric patients 13 to 17 years of age weighing ≥ 40 kg with severely active Crohn's disease and/or who have had an inadequate response or were intolerant to conventional therapy (a corticosteroid and/or aminosalicylate and/or an immunosuppressant) and/or a tumour necrosis factor alpha antagonist.
- Treatment of adult patients with chronic moderate to severe psoriasis (Ps) who are candidates for systemic therapy. For patients with chronic moderate plaque psoriasis, HUMIRA should be used after phototherapy has been shown to be ineffective or inappropriate.

Clinical use

Safety and effectiveness in pediatric patients with polyarticular JIA less than 4 years of age have not been established. Limited data are available for treatment with HUMIRA in children weighing < 15 kg. Clinical trial data for patients aged 4 to 6 years with polyarticular JIA are limited.

The safety and efficacy of HUMIRA were authorised in pediatric patients 13 to 17 years of age weighing ≥ 40 kg with severely active Crohn's disease and/or who have had an inadequate response or were intolerant to conventional therapy.

Contraindications

- Severe infections such as sepsis, tuberculosis and opportunistic infections.
- Moderate to severe heart failure (NYHA class III/IV).

Most serious warnings and precautions

Hepatosplenic T-Cell Lymphoma (HSTCL): Very rare post-marketing reports of HSTCL, a rare aggressive lymphoma that is often fatal, have been reported. Most of the patients had prior infliximab therapy as well as concomitant azathioprine or 6-mercaptopurine use for

Crohn's disease. The potential risk with the combination of azathioprine or 6-mercaptopurine and HUMIRA should be carefully considered.

Infections: Serious infections have been reported. Hospitalization or fatal outcomes associated with infections have been reported. Many of the serious infections have occurred in patients on concomitant immunosuppressive therapy that, in addition to their underlying disease, could predispose them to infections. Treatment with HUMIRA should not be initiated in patients with active infections. In patients who have been exposed to tuberculosis, and patients who have traveled in areas of high risk of tuberculosis or endemic mycoses, the risks and benefits of treatment with HUMIRA should be considered prior to initiating therapy. As with other TNF blockers, patients should be monitored closely for infections (including tuberculosis) before, during and after treatment with HUMIRA. Administration of HUMIRA should be discontinued if a patient develops a serious infection or sepsis, and appropriate therapy should be initiated. Physicians should exercise caution when considering the use of HUMIRA in patients with a history of recurrent infection or with underlying conditions which may predispose them to infections, or patients who have resided in regions where tuberculosis and histoplasmosis are endemic.

Pediatric Malignancy: Lymphoma and other malignancies, some fatal, have been reported in children and adolescent patients treated with TNF blockers, including HUMIRA.

Other relevant warnings and precautions

- Concurrent administration with other biologic DMARDs or other TNF antagonists not recommended
- Surgery: Close monitoring for infection required
- Patients with congestive heart failure: Cases of worsening congestive heart failure (CHF) and new onset CHF
- Hematologic events: Pancytopenia, including aplastic anemia, and medically significant cytopenia
- Hypersensitivity reactions, including anaphylaxis and latex allergic reactions
- Autoimmunity
- Immunosuppression
- Immunizations: Live vaccines must be avoided. It is recommended that pediatric patients, if possible, be brought up to date with all immunizations in agreement with current immunization guidelines prior to initiating HUMIRA therapy
- Infections: Tuberculosis (TB), (including reactivation and new onset of TB), opportunistic infections, (including invasive fungal infections), and hepatitis B virus reactivation
- Malignancies including lymphoma and non-lymphoma malignancy
- Neurological events: New onset or exacerbation of demyelinating disease
- Pregnant women: HUMIRA may cross the placenta; infants born to women treated with HUMIRA during pregnancy may be at increased risk for infection
- Nursing women: Breastfeeding is not recommended for at least five months after the last HUMIRA treatment
- Geriatrics: Higher incidence of infections and malignancies

For more information

Please consult the Product Monograph at <http://webprod5.hs-sc.gc.ca/dpd-bdpp/index-eng.jsp> for important information relating to adverse reactions, drug interactions, and dosing information which have not been discussed in this piece.

The Product Monograph is also available by calling at 1-888-703-3006.

† RA: rheumatoid arthritis; JIA: polyarticular juvenile idiopathic arthritis; AS: ankylosing spondylitis; PsA: psoriatic arthritis; CD: Crohn's disease; Ped CD: pediatric Crohn's disease; Ps: psoriasis

References: 1. Calin A, Garrett S, Whitlock H *et al.* A new approach to defining functional ability in ankylosing spondylitis: The development of the Bath Ankylosing Functional Index. *J Rheumatol* 1994;21:2281-85. 2. Data on file. AbbVie Corporation. 3. HUMIRA Product Monograph. AbbVie Corporation. August 21, 2013.

This form should be filled out with a nurse.

HUMIRA[®]
adalimumab
POWER TO FIGHT AS

©AbbVie Corporation
Printed in Canada
HUM/2387A01—October 2013

Member
R&D PAAB

www.abbvie.ca
1-888-703-3006



abbvie